



EUROPEAN COLORECTAL CONGRESS

Spotlight on the colon

1 – 5 December 2019, St.Gallen, Switzerland

Spotlight on the colon

Inflammatory bowel diseases (IBD)

Robotics in colorectal surgery

New technologies

Lars Pählman lecture

EAES Presidential lecture

Masterclass in colorectal surgery



Information & Registration

www.colorectalsurgery.eu


The publication of this advertisement does not constitute endorsement by the society, publisher, or Editors, and is unrelated to the content that follows



Reducing surgical mortality in Scotland by use of the WHO Surgical Safety Checklist

G. Ramsay^{1,2} , A. B. Haynes^{5,6,7}, S. R. Lipsitz⁶, I. Solsky⁵, J. Leitch⁴, A. A. Gawande^{5,8,9,10} and M. Kumar^{2,3} 

¹The Rowett Institute, University of Aberdeen, and ²Department of General Surgery, Aberdeen Royal Infirmary, Aberdeen, and ³Scottish Mortality and Morbidity Programme, Healthcare Improvement Scotland, and ⁴Healthcare Quality and Strategy, The Scottish Government, Edinburgh, UK, ⁵Safe Surgery Program, Ariadne Labs, ⁶Department of Surgical Oncology, Massachusetts General Hospital, ⁷Center for Surgery and Public Health, and ⁸Division of General and Gastrointestinal Surgery, Brigham and Women's Hospital, ⁹Department of Surgery, Harvard Medical School, and ¹⁰Department of Health Policy and Management, Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA

Correspondence to: Mr M. Kumar, Aberdeen Royal Infirmary, Foresterhill, Aberdeen AB25 2ZN, UK (e-mail: m.kumar3@nhs.net; @Manoj_K_Kumar; @online_his)

Background: The WHO Surgical Safety Checklist has been implemented widely since its launch in 2008. It was introduced in Scotland as part of the Scottish Patient Safety Programme (SPSP) between 2008 and 2010, and is now integral to surgical practice. Its influence on outcomes, when analysed at a population level, remains unclear.

Methods: This was a population cohort study. All admissions to any acute hospital in Scotland between 2000 and 2014 were included. Standardized differences were used to estimate the balance of demographics over time, after which interrupted time-series (segmented regression) analyses were performed. Data were obtained from the Information Services Division, Scotland.

Results: There were 12 667 926 hospital admissions, of which 6 839 736 had a surgical procedure. Amongst the surgical cohort, the inpatient mortality rate in 2000 was 0.76 (95 per cent c.i. 0.68 to 0.84) per cent, and in 2014 it was 0.46 (0.42 to 0.50) per cent. The checklist was associated with a 36.6 (95 per cent c.i. -55.2 to -17.9) per cent relative reduction in mortality ($P < 0.001$). Mortality rates before implementation were decreasing by 0.003 (95 per cent c.i. -0.017 to +0.012) per cent per year; annual decreases of 0.069 (-0.092 to -0.046) per cent were seen during, and 0.019 (-0.038 to +0.001) per cent after, implementation. No such improvement trends were seen in the non-surgical cohort over this time frame.

Conclusion: Since the implementation of the checklist, as part of an overall national safety strategy, there has been a reduction in perioperative mortality.

Paper accepted 4 February 2019

Published online in Wiley Online Library (www.bjs.co.uk). DOI: 10.1002/bjs.11151

Introduction

Surgery continues to be an important treatment for a wide variety of conditions with an estimated 312.9 million operations performed per year across the globe^{1,2}. Every surgical procedure has an associated risk of morbidity and death³. Multiple complex factors influence surgical outcome, with both technical and non-technical components being key factors. Consequently, surgical outcomes are influenced by multiple team members, and the systems of care in which they work^{4,5}. The rates of adverse events vary across hospitals, regions and countries, with up to half due to provider or system-wide shortcomings⁶⁻⁹. As a result, various measures to improve surgical team

performance and, thus, mitigate against surgical complications or adverse events have been advocated¹⁰.

The WHO Surgical Safety Checklist is one such measure that has been implemented internationally³. This checklist was launched in 2008 and has since become an integral part of the surgical process across the globe¹¹⁻¹³. Its aim is to make surgical procedures safer, by ensuring adherence to established practices and creating a culture of communication and teamwork that supports patient safety. The checklist is used by the entire operative team at three key points during any intervention in which harm could ensue^{3,14}. The aim of its implementation in Scotland was to improve safety of surgical procedures³, thereby improving patient

outcomes¹⁵ and mortality rates¹⁶. However, its impact on improvements in mortality rates after mandatory nationwide implementation has been questioned^{14,17–19}.

The initial pilot study²⁰ analysing the impact of checklist implementation in eight hospitals in eight separate countries noted a significant reduction in perioperative mortality rates and inpatient complications. This study had a trial format, however, and used data collated prospectively. Whether this reduction is replicated in the real-world scenario outside the context of a trial has yet to be proven. Studies^{11,12,17,21–24} performed in various health-care facilities and regions have had mixed results regarding the influence of the checklist on outcome. Implementation of the checklist by mandate in Ontario, with limited training and support, demonstrated no significant reduction in death 3 months after hospitals certified compliance¹⁷. However, in a voluntary programme in South Carolina, hospitals completing a collaborative, unit-based implementation protocol supported by educational programmes achieved a 22 per cent reduction in mortality rates²⁵. The aim of the present study was to seek to understand whether similar improvements have been seen in hospitals across Scotland, where implementation of the checklist was mandated through a national collaborative programme to improve safety of hospital healthcare services.

The Scottish Patient Safety Programme (SPSP) is a national initiative which aims to improve in-hospital mortality rates in Scotland²⁶. The SPSP was established in 2008 and had perioperative management, including implementation of the WHO surgical checklist, as one of its four key initial workstreams. This programme offered a systematic approach to improve patient safety across all hospitals in the country²⁶. It used a multidisciplinary team approach to implement key drivers for change. By implementing the surgical checklist through the perioperative arm of SPSP, Scotland coordinated its introduction nationally, with regional and local support. This study aimed to determine whether implementation of the surgical checklist, through a national-level improvement strategy combined with a unit-based partnership, had an impact on population outcomes after surgery in Scotland.

Methods

This was a population cohort study. The Information Services Division (ISD) of National Health Service Scotland collects data prospectively on all components of health service provision in the country. It uses a unique identifier which can track individual patient outcomes through time²⁷. All admissions to an acute hospital in

Scotland from 2000 to 2014 were included. Patients who were admitted to a psychiatric department, rehabilitation facility or long-term care hospital were excluded from the study.

Data were based on a continuous episode of treatment in hospital. Data were summarized and anonymized at source in ISD. Data were obtained on age, sex, deprivation, specialty, rates of return to theatre, operative urgency, cause of death and number of in-hospital deaths.

ISD follows the principles of the Data Protection Act 2018, UK²⁸. This project was reviewed by a proportionate Research Ethical Review with the Integrated Research Application System and was approved (reference number 196391). It was an observational epidemiological study and, as such, no patient involvement was possible.

Cohort definition and endpoints

The inclusion criterion was any admission to an acute-care hospital in Scotland. The surgical cohort was defined by any inpatient admission in which an operation (by OPCS-4 code association with the admission) was done between 1 January 2000 and 31 December 2014. The non-surgical cohort included all patients admitted to the same hospital settings in which no operation was undertaken. The primary endpoint was in-hospital death. The secondary endpoint in the surgical cohort was return to theatre after procedures undertaken electively.

Scottish Patient Safety Programme implementation of surgical checklist

In brief, the development was based on the Institute for Healthcare Improvement collaborative model²⁹. All local health boards were recruited to encourage clinicians, in a multidisciplinary team approach, to change the ethos around patient safety. SPSP implementations are driven nationally, but there are minor local variations to adapt to relevant specialties, context and regions.

Implementation was introduced by use of the plan–do–study–act improvement cycle. Staff education or training to support the theatre leadership to implement change was provided, and biannual nationwide meetings were held to share effective approaches to checklist implementation. These were supported by continual data collection on adherence²⁶.

The checklist was established in Scotland between 2008 and 2010. By the end of 2010, its use was nearly uniform across the country^{30,31}. The time taken for adopting the checklist appeared to be 3 years (2008–2010), so the years until 2007 were defined as the preimplementation

	Demographic information			Standardized difference (%)§		
	Preimplementation*	Implementation†	Postimplementation‡	Preimplementation versus implementation	Preimplementation versus postimplementation	Implementation versus postimplementation
	(n = 3 629 602)	(n = 1 384 425)	(n = 1 825 709)			
Women	1 962 618 (54.1)	755 201 (54.5)	985 197 (54.0)	-0.96	0.22	1.18
Age (years)						
0–19	413 402 (11.4)	137 549 (9.9)	167 837 (9.2)	4.71	7.23	2.52
20–49	1 250 792 (34.5)	442 432 (32.0)	543 558 (29.8)	5.32	10.05	4.73
50–79	1 645 698 (45.3)	669 554 (48.4)	927 711 (50.8)	-6.06	-10.97	-4.90
≥ 80	319 710 (8.8)	134 890 (9.7)	186 602 (10.2)	-3.22	-4.82	-1.59
Specialty						
General surgery	1 246 081 (34.3)	439 280 (31.7)	596 410 (32.7)	5.53	3.53	-2.01
Orthopaedics	553 391 (15.3)	242 323 (17.5)	321 701 (17.6)	-6.10	-6.41	-0.31
Other	1 830 130 (50.4)	702 822 (50.8)	907 598 (49.7)	-0.69	1.42	2.11
SIMD						
SIMD 1	882 743 (24.5)	316 824 (23.0)	409 965 (22.6)	3.45	4.50	1.05
SIMD 2	805 597 (22.3)	299 106 (21.7)	387 959 (21.3)	1.48	2.37	0.89
SIMD 3	713 275 (19.8)	281 949 (20.5)	370 022 (20.4)	-1.74	-1.48	0.26
SIMD 4	632 085 (17.5)	253 796 (18.4)	342 368 (18.8)	-2.36	-3.42	-1.07
SIMD 5	574 266 (15.9)	225 861 (16.4)	306 996 (16.9)	-1.30	-2.64	-1.33
Missing	21 626	6889	8399			
Non-elective admission	856 453 (23.6)	259 751 (18.8)	317 758 (17.4)	11.85	15.38	3.53

Values in parentheses are percentages. *Preimplementation: 2000–2001 to 2007–2008; †implementation: 2008–2009 to 2010–2011; ‡postimplementation: 2011–2012 to 2014–2015. §Standardized differences with absolute values of less than 10 per cent reflect well balanced co-variables³². SIMD, Scottish Index of Multiple Deprivation.

interval; 2008–2010 was defined as implementation; and 2011 onwards as the postimplementation interval.

Statistical analysis



Patient demographics in the three time intervals were described by use of proportions. Standardized differences were used to estimate whether these demographics were balanced over time (and thus unlikely to be confounders). Standardized differences of less than 10 per cent have been found to reflect well balanced demographics over time³². Interrupted time-series (segmented regression) analyses were used to determine whether there were changes in the level and slope of the rates during the three intervals, and not a constant downward secular trend that continued over time³³. The levels and slopes of the rates in the three time intervals were estimated with optimal weighted least squares^{34,35}, with a robust standard error to account for possible overdispersion due to clustering of outcomes within hospitals (even though hospital-level data were not available, this robust standard error can account for overdispersion)³⁶. All analyses were performed in SAS[®] version 9.4 (SAS Institute, Cary, North Carolina, USA).

Results

In Scottish hospitals between 2000 and 2014, there were 12 667 926 admissions, of which 6 839 736 had a surgical procedure. *Table 1* summarizes the demographic information for these surgical admissions. Over the preimplementation interval (2000–2001 to 2007–2008), during implementation (2008–2009 to 2010–2011) and over the postimplementation interval (2011–2012 to 2014–2015), the following co-variables were found to be well balanced with mean absolute standardized differences of less than 10 per cent: sex (female: 0.8 per cent), older age (older than 80 years: 3.2 per cent), Scottish Index of Multiple Deprivation (SIMD) (SIMD 1: 3.0 per cent) and surgical specialty (trauma and orthopaedic surgery: 4.3 per cent). Admission type was found to have a mean standardized difference greater than 10 per cent across the three intervals (non-elective admission: 10.3 per cent), indicating that this co-variable could be a potential confounder.


Overall mortality trends in surgical cohort

Fig. 1 shows the results of the time-series analysis for overall mortality rate across the three intervals. In the

Fig. 1 Time-series analysis for overall mortality rates across preimplementation, implementation and postimplementation intervals**Fig. 2 Time-series analysis for overall mortality rates adjusted for admission type across preimplementation, implementation and postimplementation intervals**

preimplementation interval, the mortality rate had an absolute decrease of 0.003 (95 per cent c.i. -0.017 to $+0.012$) per cent per year; during implementation, the annual mortality rate decreased 0.069 (-0.092 to -0.046) per cent; and in the postimplementation interval, it decreased 0.019 (-0.038 to $+0.001$) per cent. The downward trend in mortality rate seen in this model during implementation was found to be significantly different from the trend in the preimplementation interval ($P < 0.001$). Although the mortality rate trend in the postimplementation interval was not found to be significantly different from that in the preimplementation interval ($P = 0.153$), the estimate for overall mortality rate in the preimplementation interval and that in the postimplementation interval were significantly different ($P < 0.001$). Among hospital admissions with an operation performed, the inpatient mortality rate in 2000 was 0.76 (95 per cent c.i. 0.68 to 0.84) per cent and in 2014 it was 0.46 (0.42 to 0.5) per cent; if the trend in the preimplementation interval had persisted, the 2014 mortality rate would have been 0.72 (0.59 to 0.85) per cent. Thus, the intervention was associated with reduced mortality rates, demonstrating a 36.6 (95 per cent c.i. -55.2 to -17.9) per cent relative reduction over the time studied ($P < 0.001$).

A time-series analysis was also performed adjusting for admission type, which was found, on exploration of the demographic information, to be a potential confounder. The results of the adjusted model are shown in Fig. 2. Trends across the three time intervals were found to be similar in this model to those in the model for overall mortality rates shown in Fig. 1; this demonstrated that the intervention was associated with reduced mortality rates.

Fig. 3 Time-series analysis for return-to-theatre rates after admission for elective surgery across preimplementation, implementation and postimplementation intervals

Return to theatre

Fig. 3 shows the time-series analysis for return-to-theatre rate after elective admission for surgery across the three time intervals. In the preimplementation interval, the return-to-theatre rate increased 0.002 (95 per cent c.i. 0.001 to 0.002) per cent per year; during implementation, the annual return-to-theatre rate decreased 0.003 (-0.005 to -0.002) per cent; and in the postimplementation interval, it decreased 0.002 (-0.002 to -0.001) per cent. The difference between the during-implementation trend and the preimplementation trend was statistically significant ($P < 0.001$), as was that between the preimplementation

trend and the postimplementation trend ($P < 0.001$). The differences in estimates for return-to-theatre rates preimplementation compared with postimplementation were statistically significant ($P < 0.001$).

Overall mortality trends in non-surgical cohort

A time-series analysis for the 5 828 190 patients in the non-surgical cohort, with the same time frames for preimplementation, implementation and postimplementation, showed no improvement in mortality rates ($P = 0.418$).

Discussion

This nationwide population-based cohort study found a substantial reduction in mortality rates in patients undergoing surgical intervention in Scotland after the introduction and implementation of the WHO surgical checklist as part of a nationwide patient safety initiative. This reduction persisted when the data were controlled for urgency of admission, the only baseline demographic that altered during the time frame. No such trend in improvement was observed in the non-surgical cohort. From these data, it can be inferred that the improvement seen in the surgical group was associated temporally with the implementation of the surgical checklist.

The SPSP is a unique national programme that aims to change the healthcare culture to one that has patient safety at its forefront²⁶. A key element has been the testing and application of evidence-based interventions and quality-improvement methodology. These implementations have been supported locally through co-production, educational programmes and prospective data on adherence. Since its launch, the programme has contributed to a significant reduction in harm through relevant quality-improvement strategies^{26,37}. After testing, review and feedback from health boards across Scotland, the surgical checklist was included as one of the ten Patient Safety Essentials to be implemented across all health boards in Scotland³⁷. The surgical checklist was not a stand-alone intervention. It was, however, the only Patient Safety Essential that targeted surgical patients specifically during the interval studied³⁷. Thus, the addition of the checklist to the other parameters within the SPSP may have contributed to the improvement in results observed in the present study.

The implementation of each of the SPSP interventions was mandated, but with emphasis placed on collaborative working with frontline clinicians and teams, together with local clinical leadership. The SPSP ensures that healthcare improvement implementations are adopted in a nationally coordinated approach and, as such, allows patient safety

initiatives to be developed and rolled out regardless of hospital location, clinician experience or underlying knowledge. Neither checklist fidelity³⁸, nor the effectiveness of the educational programmes³⁹, has been assessed in this study. Furthermore, the surgical checklist in Scotland has predominantly involved only the clinical team, with variation noted regarding patient involvement, another area in which checklist quality could be improved⁴⁰.

Use of an observational approach at the population level reduces the risks of observation bias. The present data highlight real-world improvements outwith the context of clinical trials or research centres. There are limitations, however, to both the data and the analysis. One of the key markers would have been to assess the rates of specific events, such as wrong-site surgery. There was no access to these data in the present study; the rates of these events are so low anyway that identifying significant trends is challenging. Specific details on how each unit conducted training sessions or developed frameworks during the implementation stage were not available. However, all regions supported checklist implementation with additional structures and training of clinical staff in all theatre settings, tailored to suit each hospital and specialty.

It is acknowledged that attributing causal links to the findings in population-wide data set analysis is not possible. The data were obtained in a summarized manner and individual patient-level data were not available, precluding multivariable analysis. Thus, standardized differences, reported previously in observational research³², were used to analyse changes in demographic characteristics over time and incorporated into bivariable analysis for the effect of urgent admission, the sole factor with a mean standardized difference greater than 10 per cent.

The present findings of improved outcome are in agreement with several studies^{20,21,23,24} looking specifically at the implementation of a surgical safety checklist, but are at odds with another¹⁷ analysing population-based outcomes in the early phase after checklist initiation. The present study used a more longitudinal approach, allowing for bedding down as the checklist has become established as part of the workload culture of surgical theatre life in Scotland. This study provides further evidence that the success of checklist implementation is more pronounced when it is supported by a cohesive and wider approach to patient safety.

Acknowledgements

R. Munro and R. Black of NHS National Services Scotland, UK, provided data intelligence. B. Robson helped review this paper; A. Longmate was involved in the preliminary design of this study; J. Ingram, J. Ferbrache

and SPSP managers from health boards in Scotland provided some of the information on surgical checklist implementation practices at hospitals across NHS Scotland. Data, analytical methods and study materials used may be made available to other researchers on request. The lead author affirms that the manuscript is an honest, accurate and transparent account of the study being reported. No important aspects of the study have been omitted. Any discrepancies from the study as planned (and, if relevant, registered) have been explained. This study was not funded by any individual or group. The Research Governance Department of the University of Aberdeen sponsored this project and supported the application through ethical review and data management.

Disclosure: A.A.G. has received royalties from multiple publishers for writing on improving healthcare, including through use of checklists: Objetiva, Sextante (Brazil); Profile Books Ltd (British Commonwealth); Cheers Publishing Company, Commonwealth Publishing Co (People's Republic of China); Jesenski & Turk, Mizaik Knjiga, Mozaik Knjiga (Croatia); Dokoran (Czech Republic); Lindhart og Rinhof Forlag (Denmark); Pilgrim Group (Estonia); Editions Moyen-Courrier, Fayard, Libraire Arthème Fayard, Moyen-Courrier (France); Radarami (Georgia), S Fischer, Verlagsgruppe Random House (Germany); Crete University Press (Greece); Tericam Kindo (Hungary); Mehta Publishing House, Penguin Random House Books (India); Gramedia Pustaka Utama, Serambi Ilmu Semesta (Indonesia); Arjmand Press (Iran); Am Oved, Modan (Israel); Einaudi Editore (Italy); Misuzu Shobo, Shinyusha Co Ltd (Japan); Janis Roze Publishers (Latvia); Vaga (Lithuania); Mime Forlag (Norway); Magnum, Znak (Poland); Lua de Papel (Portugal); Codecs, Grup Media Litera, Litera, Streamland Ltd (Romania); Slovant Publishers (Slovakia); Mladinska Knjiga (Slovenia); Antoni Bosch Editor, Editorial Empuries, Galaxia Gutenberg (Spain); Asa Editore, Bookie Publishing House, Book21, Sosoh Publishing (South Korea); Volante (Sweden); Alpina, AST (Russia); Matchon, Openworlds (Thailand); Arbeiderspers, Nieuwezijds, Uitgeverij De Arbeiderspers, Uitgeverij Nieuwezijds (the Netherlands); Domingo, Koton Kitap (Turkey); Verlagsgruppe Vivat (Ukraine); CBS Television, Henry Holt, Houghton Mifflin, Harvard Business School Press, McGraw Hill, Pearson Publishing, Public Broadcasting Service, Picador USA (USA), First News-Tn Viet Publishing Co, Suc Manh Ngoi But Co (Vietnam); and Harper Collins (World). The authors declare no other conflict of interest.

References

- 1 Shrimme MG, Bickler SW, Alkire BC, Mock C. Global burden of surgical disease: an estimation from the provider perspective. *Lancet Glob Health* 2015; 3(Suppl 2): S8–S9.
- 2 Weiser TG, Haynes AB, Molina G, Lipsitz SR, Esquivel MM, Uribe-Leitz T *et al.* Size and distribution of the global volume of surgery in 2012. *Bull World Health Organ* 2016; **94**: 201–209F.
- 3 WHO. *WHO Guidelines for Safe Surgery 2009: Safe Surgery Saves Lives*. WHO, Geneva; 2009.
- 4 Youngson GG. Nontechnical skills in pediatric surgery: factors influencing operative performance. *J Pediatr Surg* 2016; **51**: 226–230.
- 5 Sacks GD, Shannon EM, Dawes AJ, Rollo JC, Nguyen DK, Russell MM *et al.* Teamwork, communication and safety climate: a systematic review of interventions to improve surgical culture. *BMJ Qual Saf* 2015; **24**: 458–467.
- 6 GlobalSurg Collaborative. Mortality of emergency abdominal surgery in high-, middle- and low-income countries. *Br J Surg* 2016; **103**: 971–988.
- 7 Healey MA, Shackford SR, Osler TM, Rogers FB, Burns E. Complications in surgical patients. *Arch Surg* 2002; **137**: 611–617.
- 8 Gawande AA, Thomas EJ, Zinner MJ, Brennan TA. The incidence and nature of surgical adverse events in Colorado and Utah in 1992. *Surgery* 1999; **126**: 66–75.
- 9 Kable AK, Gibberd RW, Spigelman AD. Adverse events in surgical patients in Australia. *Int J Qual Health Care* 2002; **14**: 269–276.
- 10 Maggard-Gibbons M. The use of report cards and outcome measurements to improve the safety of surgical care: the American College of Surgeons National Surgical Quality Improvement Program. *BMJ Qual Saf* 2014; **23**: 589–599.
- 11 Lilaonitkul M, Kwikiriza A, Ttendo S, Kiwanuka J, Munyarungero E, Walker IA *et al.* Implementation of the WHO Surgical Safety Checklist and surgical swab and instrument counts at a regional referral hospital in Uganda – a quality improvement project. *Anaesthesia* 2015; **70**: 1345–1355.
- 12 Epiu I, Tindimwebwa JVB, Mijumbi C, Ndarugirire F, Twagirumugabe T, Lugazia ER *et al.* Working towards safer surgery in Africa: a survey of utilization of the WHO safe surgical checklist at the main referral hospitals in East Africa. *BMC Anesthesiol* 2016; **16**: 60.
- 13 de Jager E, McKenna C, Bartlett L, Gunnarsson R, Ho Y-H. Postoperative adverse events inconsistently improved by the World Health Organization Surgical Safety Checklist: a systematic literature review of 25 studies. *World J Surg* 2016; **40**: 1842–1858.
- 14 Walker IA, Reshamwalla S, Wilson IH. Surgical safety checklists: do they improve outcomes? *Br J Anaesth* 2012; **109**: 47–54.
- 15 Haugen AS, Wæhle HV, Almeland SK, Harthug S, Sevdalis N, Eide GE *et al.* Causal analysis of World Health Organization's surgical safety checklist implementation quality and impact on care processes and patient outcomes: secondary analysis from a large stepped wedge cluster

- randomized controlled trial in Norway. *Ann Surg* 2019; **269**: 283–290.
- 16 Weiser TG, Haynes AB. Ten years of the Surgical Safety Checklist. *Br J Surg* 2018; **105**: 927–929.
 - 17 Urbach DR, Govindarajan A, Saskin R, Wilton AS, Baxter NN. Introduction of surgical safety checklists in Ontario, Canada. *N Engl J Med* 2014; **370**: 1029–1038.
 - 18 Dharampal N, Cameron C, Dixon E, Ghali W, Quan ML. Attitudes and beliefs about the surgical safety checklist: just another tick box? *Can J Surg* 2016; **59**: 268–275.
 - 19 Mascherek AC, Bezzola P, Gehring K, Schwappach DLB. [Effect of a two-year national quality improvement program on surgical checklist implementation.] *Z Evid Fortbild Qual Gesundheitswes* 2016; **114**: 39–47.
 - 20 Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat A-HS, Dellinger EP *et al*. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med* 2009; **360**: 491–499.
 - 21 Mayer EK, Sevdalis N, Rout S, Caris J, Russ S, Mansell J *et al*. Surgical Checklist Implementation Project: the impact of variable WHO checklist compliance on risk-adjusted clinical outcomes after national implementation: a longitudinal study. *Ann Surg* 2016; **263**: 58–63.
 - 22 de Vries EN, Prins HA, Crolla RMPH, den Outer AJ, van Andel G, van Helden SH *et al*. Effect of a comprehensive surgical safety system on patient outcomes. *N Engl J Med* 2010; **363**: 1928–1937.
 - 23 Bock M, Fanolla A, Segur-Cabanac I, Auricchio F, Melani C, Girardi F *et al*. A comparative effectiveness analysis of the implementation of surgical safety checklists in a tertiary care hospital. *JAMA Surg* 2016; **151**: 639–646.
 - 24 Abbott TEF, Ahmad T, Phull MK, Fowler AJ, Hewson R, Biccard BM *et al*. The surgical safety checklist and patient outcomes after surgery: a prospective observational cohort study, systematic review and meta-analysis. *Br J Anaesth* 2018; **120**: 146–155.
 - 25 Haynes AB, Edmondson L, Lipsitz SR, Molina G, Neville BA, Singer SJ *et al*. Mortality trends after a voluntary checklist-based surgical safety collaborative. *Ann Surg* 2017; **266**: 923–929.
 - 26 Haraden C, Leitch J. Scotland's successful national approach to improving patient safety in acute care. *Health Aff (Millwood)* 2011; **30**: 755–763.
 - 27 Womersley J. The public health uses of the Scottish Community Health Index (CHI). *J Public Health Med* 1996; **18**: 465–472.
 - 28 UK Parliament. *Data Protection Act 2018 (c12)*. The Stationery Office: London, 2018.
 - 29 Institute for Healthcare Improvement. The Breakthrough Series: IHI's collaborative model for achieving breakthrough improvement. *Diabetes Spectrum* 2004; **17**: 97–101.
 - 30 Freeman T. *Special Report: Leading on Safety*; 2014. <https://www.holyrood.com/articles/special-report/special-report-leading-safety> [accessed 1 June 2018].
 - 31 Foster A. *Highlights of the Scottish Patient Safety Programme National Conference – Driving Improvements in Patient Safety*; 2014. http://www.healthcareimprovementscotland.org/our-work/patient_safety/programme_resources/spsp_conference_report.aspx [accessed 8 March 2018].
 - 32 Austin PC. Using the standardized difference to compare the prevalence of a binary variable between two groups in observational research. *Commun Stat Simul Comput* 2009; **38**: 1228–1234.
 - 33 Penfold RB, Zhang F. Use of interrupted time series analysis in evaluating health care quality improvements. *Acad Pediatr* 2013; **13**: S38–S44.
 - 34 Grizzle JE, Starmer CF, Koch GG. Analysis of categorical data by linear models. *Biometrics* 1969; **25**: 489–504.
 - 35 Lipsitz SR. Methods for estimating the parameters of a linear model for ordered categorical data. *Biometrics* 1992; **48**: 271–281.
 - 36 White H. Maximum likelihood estimation of misspecified models. *Econometrica* 1982; **50**(1): 1–25.
 - 37 Leitch J. *Next Steps for Acute Adult Safety – Patient Safety Essentials and Safety Priorities*. CEL19; 2013. https://www.sehd.scot.nhs.uk/mels/cel2013_19.pdf [accessed 1 June 2018].
 - 38 Levy SM, Senter CE, Hawkins RB, Zhao JY, Doody K, Kao LS *et al*. Implementing a surgical checklist: more than checking a box. *Surgery* 2012; **152**: 331–336.
 - 39 Flin R, Youngson G, Yule S, Paterson-Brown S, Moran N. *The Non-Technical Skills for Surgeons (NOTSS) System Handbook v1.2. Structuring Observation, Rating and Feedback of Surgeons' Behaviours in the Operating Theatre*. University of Aberdeen: Aberdeen, 2012; 1–18.
 - 40 Russ SJ, Rout S, Caris J, Moorthy K, Mayer E, Darzi A *et al*. The WHO surgical safety checklist: survey of patients' views. *BMJ Qual Saf* 2014; **23**: 939–946.